著書・論文

著者	著書	論文名	出版書店名	掲載誌名	掲載巻	掲載ページ
Yoshizawa N, Sugimoto K, Tameda M, Inagaki Y, Ikejiri M, Inoue H, Usui M, Ito M, Takei Y.		miR-3940-5p/miR-8069 ratio in urine exosomes is a novel diagnostic biomarker for pancreatic ductal adenocarcinoma.		Oncol Lett.	19(4)	2677-2684
Ito T, Kuriyama N, Kozuka Y, Komatsubara H, Ichikawa K, Noguchi D, Hayasaki A, Fujii T, Iizawa Y, Kato H, Tanemura A, Murata Y, Kishiwada M, Mizuno S, Usui M, Sakurai H, Isaji S.		High tumor budding is a strong predictor of poor prognosis in the resected perihilar cholangiocarcinoma patients regardless of neoadjuvant therapy, showing survival similar to those without resection.		BMC Cancer	20(1)	doi: 10.1186/s12885-020-6695-9
Hayasaki A, Murata Y, Usui M, Hibi T, Fujii T, Iizawa Y, Kato H, Tanemura A, Azumi Y, Kuriyama N, Kishiwada M, Mizuno S, Sakurai H, Uchida K, Isaji S.		Clinical Significance of Plasma Apolipoprotein-AII Isoforms as a Marker of Pancreatic Exocrine Disorder for Patients with Pancreatic Adenocarcinoma Undergoing Chemoradiotherapy, Paying Attention to Pancreatic Morphological Changes.		Biomed Res Int.	第4回	doi: 10.1155/2019/5738614. eCollection 2019.
Ogura S, Tameda M, Sugimoto K, Ikejiri M, Usui M, Ito M, Takei Y.		A substitution in the pre-S1 promoter region is associated with the viral regulation of hepatitis B virus.		Virol J.	16(1)	doi: 10.1186/s12985-019-1169-x.
Ichikawa K, Mizuno S, Hayasaki A, Kishiwada M, Fujii T, Iizawa Y, Kato H, Tanemura A, Murata Y, Azumi Y, Kuriyama N, Usui M, Sakurai H, Isaji S.		IPrognostic Nutritional Index After Chemoradiotherapy Was the Strongest Prognostic Predictor Among Biological and Conditional Factors in Localized Pancreatic Ductal Adenocarcinoma Patients.		Cancers	11(4)	doi: 10.3390/cancers11040514.
臼井正信、東口髙志		食事摂取・栄養状態のアセスメント-低栄養の新たな診断基準GLIM criteriaを中心に一		「看護技術」特集企画：がん治療中の『食』を考		
臼井正信、東口髙志		免疫低下症例に対するプロバイオティクス，特に酪酸菌の重要性		雲出川出版		
伊藤彰博、東口髙志、臼井正信、都築則正、二村昭彦		がん緩和ケアにおける栄養管理		Progress in Medicine	39(8)	811-817